

**Response of leaf and fine roots proteomes of *Salix viminalis* L. to growth on Cr-rich tannery waste**

*Environmental Science and Pollution Research*

Agata Zemleduch-Barylska\*, Gabriela Lorenc-Plucińska


Institute of Dendrology Polish Academy of Sciences, Parkowa 5, 62-035 Kórnik, Poland

\*Corresponding author: Agata Zemleduch-Barylska

Tel.: +48 697418025; fax: +48 618170166

E-mail address: [agata.zemleduch@wp.pl](mailto:agata.zemleduch@wp.pl)

**Supplementary Fig. 1.** Representative 2-DE gel obtained for the **leaves** of *S. viminalis* grown on a control soil or tannery waste. Differentially abundant protein spots (at least twofold variation at  $p < 0.05$ ) identified with MS analysis are marked with arrows. Histograms show the distribution of relative spots volumes in each growth conditions (a = tannery waste, b = control soil).



**Supplementary Fig. 2.** Representative 2-DE gel obtained for the **fine roots** of *S. viminalis* grown on a control soil or tannery waste. Differentially abundant protein spots (at least twofold variation at  $p < 0.05$ ) identified with MS analysis are marked with arrows. Histograms show the distribution of relative spots volumes in each growth conditions (a = tannery waste, b = control soil).

